

**SIXTY-FIRST ANNUAL
WILLIAM LOWELL PUTNAM MATHEMATICAL COMPETITION**

Saturday, December 2, 2000

Examination A

A1. Let A be a positive real number. What are the possible values of $\sum_{j=0}^{\infty} x_j^2$, given that $x_0, x_1, x_2, \dots$ are positive numbers for which $\sum_{j=0}^{\infty} x_j = A$?

A2. Prove that there exist infinitely many integers n such that n , $n + 1$, and $n + 2$ are each the sum of two squares of integers.

[*Example:* $0 = 0^2 + 0^2$, $1 = 0^2 + 1^1$, and $2 = 1^1 + 1^1$.]

A3. The octagon $P_1P_2P_3P_4P_5P_6P_7P_8$ is inscribed in a circle, with the vertices around the circumference in the given order. Given that the polygon $P_1P_3P_5P_7$ is a square of area 5 and the polygon $P_2P_4P_6P_8$ is a rectangle of area 4, find the maximum possible area of the octagon.

A4. Show that the improper integral

$$\lim_{B \rightarrow \infty} \int_0^B \sin(x) \sin(x^2) dx$$

converges.

A5. Three distinct points with integer coordinates lie in the plane on a circle of radius $r > 0$. Show that two of these points are separated by a distance of at least $r^{1/3}$.

A6. Let $f(x)$ be a polynomial with integer coefficients. Define a sequence $A_0, a_1, \dots$ of integers such that $a_0 = 0$ and $a_{n+1} = f(a_n)$ for all $n \geq 0$. Prove that if there exists a positive integer m for which $a_m = 0$, then either $a_1 = 0$ or $a_2 = 0$.

**SIXTY-FIRST ANNUAL
WILLIAM LOWELL PUTNAM MATHEMATICAL COMPETITION**

Saturday, December 2, 2000

Examination B

B1. Let a_j , b_j , and c_j be integers for $1 \leq j \leq N$. Assume, for each j , that at least one of a_j, b_j, c_j is odd. Show that there exist integers r, s, t such that $ra_j + sb_j + tc_j$ is odd for at least $4N/7$ values of j , $1 \leq j \leq N$.

B2. Prove that the expression

$$\frac{\gcd(m, n)}{n} \binom{n}{m}$$

is an integer for all pairs of integers $n \geq m \geq 1$. [Here $\binom{n}{m} = \frac{n!}{m!(n-m)!}$ and $\gcd(m, n)$ is the greatest common divisor of m and n .]

B3. Let $f(t) = \sum_{j=1}^N a_j \sin(2\pi jt)$, where each a_j is real and $a_N \neq 0$. Let N_k denote the number of zeros (including multiplicities) of $\frac{d^k f}{dt^k}$. Prove that

$$N_0 \leq N_1 \leq N_2 \leq \cdots \quad \text{and} \quad \lim_{k \rightarrow \infty} N_k = 2N.$$

[Added note: Presumably, one is to restrict t to the interval $[0, 1)$ when counting the zeros.]

B4. Let $f(x)$ be a continuous function such that $f(2x^2 - 1) = 2xf(x)$ for all x . Show that $f(x) = 0$ for $-1 \leq x \leq 1$.

B5. Let S_0 be a finite set of positive integers. We define finite sets $S_1, S_2, \dots$ of positive integers as follows:

Integer a is in S_{n+1} if and only if exactly one of $a - 1$ or a is in S_n .

Show that there exist infinitely many integers N for which $S_N = S_0 \cup \{N + a : a \in S_0\}$.

B6. Let B be a set of more than $2^{n+1}/n$ distinct points with coordinates of the form $(\pm 1, \pm 1, \dots, \pm 1)$ in n -dimensional space, with $n \geq 3$. Show that there are three distinct points in B which are the vertices of an equilateral triangle.